

Satellite potentials for hypergeometric Natanzon potentials

S Codriansky^{a1} and S Salamo^{b2}

^a*Departamento de Matemáticas y Física, Instituto Pedagógico de Caracas
Universidad Pedagógica Experimental Libertador*

^b*Universidad Simón Bolívar, Departamento de Física Apartado Postal 89000,
Caracas, Venezuela*

PACS 03.65.Fd - Algebraic methods

Abstract

As a result of the $so(2,1)$ treatment of the hypergeometric Natanzon potentials a set of potentials related to a given one is determined. The set arises as a result of the action of the $so(2,1)$ generators on the carrier space of an irreducible representation.

1 Introduction

In [1] an $so(2,1)$ description of a given hypergeometric Natanzon [2] potential V_N was presented. The study of the discrete spectrum led to the result that three parameters (called group parameters) are required to completely describe the eigenstates. Two of these parameters correspond to labels of states in a particular irreducible representation (irrep) of $so(2,1)$: the eigenvalues q of the Casimir operator and m of the compact generator. The third parameter, p , labels a particular set of $so(2,1)$ generators. It is found that in most cases there is one eigenstate for each value of p and as a result different sets of $so(2,1)$ generators, each for a given (allowed) value of p , are necessary to describe the eigenfunctions of V_N . It is unusual that all the eigenstates of V_N belong to a single irrep so that if attention is fixed in a particular irrep the states connected by the $so(2,1)$ generators are, in the majority of cases, associated with different Natanzon potentials. It is this fact that suggests the definition of a satellite potential. The construction of the set of satellite potentials associated to V_N is the content of this paper. The term satellite potential is used in the present paper as in [3].

The action of the $so(2,1)$ generators defines a set of potentials which

¹codrians@reacciun.ve

²ssalamo@fis.usb.ve

share the property that the eigenstates connected by them have the same p and q but correspond, in general, to different energy eigenvalues. If the energy eigenvalues of the states connected by the $so(2,1)$ generators are the same the following question naturally arises: are the $so(2,1)$ generators related to the operators in supersymmetric quantum mechanics (SUSYQM) [4] This is not so for the shape invariant potentials V_S [5] considered in this paper, as is shown below. It is also proven that the state reached by the action of the $so(2,1)$ generators belongs to a Natanzon potential. This implies that the Natanzon class of potentials is invariant under this algebra, a result different to the one obtained in SUSYQM due to the fact that the SUSYQM operators may define a potential that does not belong to the Natanzon class [8].

2 The $so(2,1)$ description of the hypergeometric Natanzon potentials

A two variable realization of the algebra $so(2,1)$ is used with the generators taken as

$$e^{\pm i\phi} J_{\mp} = -\frac{i(1+z)}{2\sqrt{z}} \frac{\partial}{\partial \phi} \mp \frac{\sqrt{z}(z-1)}{z'} \frac{\partial}{\partial r} + \left(\mp \frac{\sqrt{z}(z-1)z''}{2z'^2} + \frac{(z-1)(\pm 1+p)}{2\sqrt{z}} \right) \quad (1)$$

where $z = z(r)$ and $z' = dz/dr$.

The expressions (1) lead to the Casimir, $Q = J_0(J_0 + 1) - J_- J_+$,

$$Q = (z-1)^2 \left[\frac{z}{z'^2} \frac{\partial^2}{\partial r^2} + \frac{1}{4z} \frac{\partial^2}{\partial \phi^2} + \frac{ip(1+z)}{2z(z-1)} \frac{\partial}{\partial \phi} + \frac{-p^2+1}{4z} + \frac{z}{2} \frac{z'''}{z'^3} - 3 \frac{z}{4} \frac{z''^2}{z'^4} \right] \quad (2)$$

The significant result that follows from (1, 2) is the appearance of the parameter p in the explicit expression for the generators and the Casimir; this constant distinguishes a particular set of $so(2,1)$ generators and plays a crucial role in the $so(2,1)$ description of the Natanzon potentials.

The physical problem dealt with is the derivation of the discrete spectrum of the Hamiltonian and to this end the compact operator J_0 is diagonalized; the considered irreps of $so(2,1)$ are unitary and therefore infinite dimensional and of these it is relevant the one bounded below (the D^+ representation). In this representation the eigenvalue m of the compact operator is given in terms of the eigenvalue q of the Casimir and the counter $\nu = 0, 1, \dots$ as

$$m(\nu) = \nu + \frac{1}{2} + \frac{1}{2} \sqrt{4q+1} \quad (3)$$

With the above results the $so(2,1)$ description of the Schrödinger equation is defined by (4), referred to as the master equation

$$G(r) (Q - q) \Psi(r, \phi) = (E - H) \Psi(r, \phi) \quad (4)$$

where H is the Hamiltonian and q, E the eigenvalues of the Casimir and Hamiltonian, respectively. The function $G(r)$ ensures that the coefficient of the second derivative of $\Psi(r, \phi)$ are the same on both sides. From (4) it follows that in general q could be a function of ν which also labels E ; in spite of the fact that p does not appear explicitly in (4) it could also depend on ν . Each eigenfunction of the Casimir (equivalently of the Hamiltonian) has the form

$$\Psi(r, \phi) = \exp(im(\nu)\phi)g \quad (5)$$

and is also an eigenfunction of the compact operator J_0 . The function $g = g(r)$ is determined solving the master equation.

The set of Natanzon hypergeometric potentials [2] is given by

$$\begin{aligned} V_N = & \frac{f z^2 - (h_0 - h_1 + f) z + h_0 + 1}{R} \\ & + \left[a + \frac{a + (c_1 - c_0)(2z - 1)}{z(z - 1)} - \frac{5\Delta}{4R} \right] \left[\frac{z(1 - z)}{R} \right]^2 \end{aligned} \quad (6)$$

where $(a, c_0, c_1, f, h_0, h_1)$ are the Natanzon parameters and z is a solution of the differential equation

$$z(r)' = 2 \frac{z(1 - z)}{\sqrt{R}}; \quad (7)$$

the other symbols that appear in (6, 7) are given by

$$\tau = c_1 - c_0 - a, \quad \Delta = \tau^2 - 4ac_0, \quad R = az^2 + \tau z + c_0 \quad (8)$$

The $so(2,1)$ description of $V_N(r)$ is obtained after the explicit expression for the Casimir (2) is put into a form similar to (6) after use of (7) and the coefficients of the powers of z compared; this leads to

$$\begin{aligned} p(\nu) + m(\nu) &= \sqrt{-aE(\nu) + f + 1} = \alpha(\nu) \\ p(\nu) - m(\nu) &= \sqrt{-c_0E(\nu) + h_0 + 1} = \beta(\nu) \\ \sqrt{4q(\nu) + 1} &= \sqrt{-c_1E(\nu) + h_1 + 1} = \delta(\nu). \end{aligned} \quad (9)$$

From (3, 9) $p(\nu), q(\nu), m(\nu)$ -called the group parameters- and $E(\nu)$ are determined for each value of ν thus fixing a particular $so(2,1)$ irrep and the energy eigenvalue. This implies that each eigenfunction belongs to the carrier space of an $so(2,1)$ which depends on p see (2) and hence may depend on ν as follows from (9). From now on $p(\nu), q(\nu)$ and $m(\nu)$ are written p, q and m . The carrier space of a specific irrep is given by

$$\Psi_{pqm}(r, \phi) = \exp(im\phi)\Phi_{pqm}(r) \quad (10)$$

where

$$\Phi_{pqm}(r) =$$

$$K z^{\beta(\nu)/2} (1-z)^{\delta(\nu)/2} R^{1/4} {}_2F_1(-\nu, \alpha(\nu) - \nu; 1 + \beta(\nu); z). \quad (11)$$

where K is a normalization constant. The energy spectrum is obtained from

$$\alpha(\nu) - \beta(\nu) - \delta(\nu) = 2\nu + 1. \quad (12)$$

3 Satellite potentials

In this Section the possibilities allowed by (9) for fixed values of the group parameters (p, q) are presented. The study of this situation leads to the construction of the satellite potentials. The action of the $so(2, 1)$ generators on (11) is the following

$$\begin{aligned} J_- \Psi_{pqm} &= \frac{\nu (\alpha(\nu) - \nu - 1 - \beta(\nu))}{1 + \beta(\nu)} \Psi_{pqm-1} \\ J_+ \Psi_{pqm} &= -\beta(\nu) \Psi_{pqm+1} \end{aligned} \quad (13)$$

which follow from [7]

$$\begin{aligned} {}_2F_1(a+1, b+1, c+1; z) &= \\ &- \frac{1}{zb(z-1)a} [c(c-1) {}_2F_1(a-1, b, c-1; z) - c(zb-c+1) {}_2F_1(a, b, c; z)] \end{aligned} \quad (14)$$

and

$$\begin{aligned} {}_2F_1(a-1, b, c-1; z) &= \\ &- \frac{1}{(c-1)} [(b-c+1) {}_2F_1(a, b, c; z) - b(z-1) {}_2F_1(a, b+1, c; z)] \end{aligned} \quad (15)$$

It follows from (13) that the action of the $so(2, 1)$ generators on an eigenfunction of a given Natanzon potential gives an eigenfunction of a different Natanzon potential. In fact, from (13) the same z is present so that a, c_0, c_1 are unchanged. The other three Natanzon parameters f_1, h_{01}, h_{11} depend on ν after (13) and (9) are determined by

$$\alpha_1(\nu \pm 1) = \alpha(\nu) \pm 1, \beta_1(\nu \pm 1) = \beta(\nu) \mp 1, \delta_1(\nu \pm 1) = \delta(\nu) \quad (16)$$

where $\alpha_1(\nu), \beta_1(\nu), \delta_1(\nu)$ follow from (9) by $f \rightarrow f_1, h_0 \rightarrow h_{01}, h_1 \rightarrow h_{11}$. Thus the complete set of Natanzon parameters is determined for the new potential which shows that the Natanzon class is invariant under the action of the $so(2, 1)$ generators. Nevertheless the energy eigenvalue $E_1(\nu)$ is undetermined since (12) is a linear combination of (9). The fact that one of the four quantities $(f, h_0, h_1, E(\nu))$ is arbitrary can be used to fix the energy of the ground state.

The potentials to which the eigenfunctions with $m \pm 1$ are associated will be called satellite to the one related to the eigenfunction labeled by m . The common feature to all these potentials is that the values of both p and q remain

unchanged. It is natural to ask, how many satellite potentials are associated to a given one? Different eigenfunctions of a given Natanzon potential do not necessarily belong to the same $so(2, 1)$ irrep and therefore for each such function a certain number of satellite potentials is determined. Fix the attention in one eigenfunction. It belongs at the same time to the set of eigenfunctions of V_N and to the $so(2, 1)$ irrep. Recall that ν labels the place of $\Phi_{pqm}(r)$ on the irrep and call λ the label of its place in the set of eigenfunctions of V_N . The above description assigns a set of values of (p, q, m) in each case and both sets must coincide, that is

$$p(\nu) = p(\lambda), q(\nu) = q(\lambda), m(\nu) = m(\lambda) \quad (17)$$

which imply

$$\lambda = \nu \quad (18)$$

so that the numerical value of both labels coincide on the irrep. Therefore, the action of J_+ increase both ν and λ and if there is a maximum value of λ for a given potential a finite number of satellite potentials will be constructed. The maximum value λ depends on whether the $r \rightarrow \infty$ limit is finite or not for each of the satellite potentials.

To complete the answer consider a Natanzon potential V_N one of whose eigenfunctions, $\Psi_{\lambda_0=\nu_0}^{\nu_0}$, is in the (p, q) irrep; its place in the set of eigenfunctions is $\lambda = \lambda_0$ and the one in the irrep is $\nu = \nu_0$. The reason for introducing both ν and λ (which may seem redundant because for this particular function their numerical values coincide) is that, as shown above, the Natanzon parameters of the satellite to V_N depend on those for V_N in a way determined by (16); therefore, these parameters may (and in fact will) be functions of ν . Thus, displacement along the irrep produces sets of Natanzon parameters that change with ν in such a way that their numerical values coincide with those for V_N when $\nu = \nu_0$ and those for the satellite when $\nu = \nu_0 \pm 1$. Moving from one eigenfunction of V_N to another keeps ν_0 fixed since in this case the Natanzon parameters do not change. The question now is the following: if $\Psi_{\lambda_0=\nu_0}^{\nu_0}$ and $\Psi_{\lambda_0+1=\nu_0+1}^{\nu_0+1}$ are in the same irrep, does the same occurs for $\Psi_{\lambda_0+1}^{\nu_0}$ and $\Psi_{\lambda_0+2}^{\nu_0+1}$? To answer this question the system (9, 3) is studied taking the Natanzon parameters for $\nu = \nu_0$ and $\nu = \nu_0 + 1$, replacing ν by λ and comparing the values of p and q obtained for $\lambda = \lambda_0 + 1$ (for ν_0) and $\lambda = \lambda_0 + 2$ (for $\nu_0 + 1$). It turns out that in general these values are not the same, the analysis described above has to be repeated for each eigenfunction and a new set of satellite potentials is thus determined.

4 Particular cases

The results that follow include the set of Natanzon parameters for the potential and its supersymmetric partner and the rule that generates the parameters for the satellite potentials. It has been found that for the six cases studied

the satellite potentials do not coincide with the SUSY partner. A detailed study along the lines presented in this paper has been performed for the Eckart potential in [6]; where it is shown that the satellite potential does not coincide with the supersymmetric partner.

1. The Pöschl-Teller II potential in the notation of [9] is

$$V_{PT2} = (A - B)^2 - A(A + \alpha) \operatorname{sech}(\alpha r)^2 + B(B - \alpha) \operatorname{csch}(\alpha r)^2. \quad (19)$$

The Natanzon parameters

$$\begin{aligned} a &= 0, \quad c_0 = 0, \quad c_1 = \alpha^{-2}, \\ f &= \frac{(2A - \alpha)(2A + 3\alpha)}{4\alpha^2}, \\ h_0 &= \frac{(2B + \alpha)(2B - 3\alpha)}{4\alpha^2}, \\ h_1 &= \frac{(A - B + \alpha)(A - B - \alpha)}{\alpha^2} \end{aligned} \quad (20)$$

reproduce V_{PT2} with $z = \tanh(\alpha r)^2$ replaced in (6). From (9) the group parameters are

$$p = \frac{A + B}{2\alpha}, \quad q = \frac{(A - B - 2\nu\alpha)^2 - \alpha^2}{4\alpha^2}, \quad m = \frac{A - B + \alpha}{2\alpha}. \quad (21)$$

and again from (9)

$$\alpha(\nu) = \frac{\alpha + 2A}{2\alpha}, \quad \beta(\nu) = -\frac{-2B + \alpha}{2\alpha}, \quad \delta(\nu) = -\frac{2\nu\alpha - A + B}{\alpha} \quad (22)$$

The energy spectrum is obtained from (12) and (22) as

$$E(\nu) = -4\alpha\nu(\nu\alpha - A + B) \quad (23)$$

After use of (16), the action of J_+ leads to

$$\alpha_1(\nu + 1) = \frac{2A + 3\alpha}{2\alpha}, \quad \beta_1(\nu + 1) = \frac{2B - 3\alpha}{2\alpha}, \quad \delta_1(\nu + 1) = 2\nu + \frac{B - A}{\alpha}; \quad (24)$$

calling (A_S, B_S) the parameters of the potential in (22) and equate the result with the values in (24) to obtain

$$A_S = A + \alpha, \quad B_S = B - \alpha \quad (25)$$

The energy spectrum for the satellite potential follows from (12)

$$E_S(\nu + 1) = E(\nu) + h_{1S}\alpha^2 + \alpha^2 + 2AB - B^2 - A^2 \quad (26)$$

with h_{1S} arbitrary; if the condition that the energy vanishes for $\nu = 0$ is imposed, then

$$h_{1S} = \frac{(-A + B - \alpha)(\alpha + B - A)}{\alpha^2}. \quad (27)$$

which leads to $E_S(\nu + 1) = E(\nu)$. Notice that, from (25), it follows that the change in the parameters does not coincide with the one found in SUSYQM to relate a potential with its supersymmetric partner [9]. The chain of satellite potentials is, thus, different from the chain of supersymmetric partners. Since the choice (27) produces equal values of the energy for both the potential and its satellite it is natural to compare this result with the one obtained in the potential algebra of [10] which consider the case $B = 0$ and no constant term. The change (25) take the value of B to $-\alpha$ and therefore to a completely different Pöschl-Teller potential; the conclusion is that the set of satellite potentials is different to the set of potentials obtained with the potential algebra.

2. The Rosen-Morse potential

$$V_{RM} = A^2 + \frac{B^2}{A^2} + 2B \tanh(\alpha r) - A(A + \alpha) \operatorname{sech}(\alpha r)^2 \quad (28)$$

is obtained from (6) with the Natanzon parameters

$$\begin{aligned} a &= 0, \quad c_0 = c_1 = 1/\alpha^2 \\ f &= 4 \frac{A(A + \alpha)}{\alpha^2}, \\ h_0 &= \frac{(-B + A\alpha + A^2)(-B - A\alpha + A^2)}{\alpha^2 A^2}, \\ h_1 &= \frac{(B + A\alpha + A^2)(B - A\alpha + A^2)}{\alpha^2 A^2} \end{aligned} \quad (29)$$

with $z = 1/2 + \tanh(\alpha r)/2$. The group parameters are found to be

$$\begin{aligned} p &= \frac{2A}{\alpha} - m + 1, \\ q &= (\nu + 1 - m)(\nu - m), \\ m &= \frac{-A^2 + \nu \alpha^2 - A\alpha - B + \nu^2 \alpha^2}{2(\nu \alpha - A)\alpha} \end{aligned} \quad (30)$$

From (9, 30) it follows

$$\begin{aligned} \alpha(\nu) &= \frac{2A + \alpha}{\alpha}, \\ \beta(\nu) &= \frac{\nu^2 \alpha^2 - 2\nu \alpha A + A^2 - B}{(A - \nu \alpha)\alpha}, \\ \delta(\nu) &= \frac{A^2 + B + \nu^2 \alpha^2 - 2\nu \alpha A}{(A - \nu \alpha)\alpha} \end{aligned} \quad (31)$$

and from (12) the energy spectrum is given by

$$E(\nu) = A^2 - (A - \nu \alpha)^2 + \frac{B^2}{A^2} - \frac{B^2}{(A - \nu \alpha)^2} \quad (32)$$

For the satellite potential it is found from (12)

$$\begin{aligned}\alpha_1(\nu+1) &= 2 \frac{A+\alpha}{\alpha}, \\ \beta_1(\nu+1) &= -\frac{A-\nu\alpha}{\alpha} - 1 + \frac{B}{(A-\nu\alpha)\alpha}, \\ \delta_1(\nu+1) &= -\frac{A-\nu\alpha}{\alpha} - \frac{B}{(A-\nu\alpha)\alpha}\end{aligned}\tag{33}$$

Setting (A_S, B_S, p_S, q_S) instead of (A, B, p, q) in (30) with $\nu \rightarrow \nu+1$ and requiring that $p(\nu) = p_S(\nu+1), q(\nu) = q_S(\nu+1)$ it is found

$$A_S = A + \alpha/2, \quad B_S = -\frac{(2\nu\alpha + \alpha - 2A)(\nu\alpha^2 - A\alpha - 2B)}{4\nu\alpha - A}\tag{34}$$

The energy of the satellite is found to be

$$E_S(\nu+1) = E(\nu) + (h_{1S} - h_1) \alpha^2\tag{35}$$

with obvious meaning for h_{1S} . The same comment made for the previous example are valid here.

The conclusion that follows from the above examples is that the chain of potentials defined by the action of the $so(2,1)$ generators is different from the one defined by the action of the SUSYQM operators and of the potential algebra.

References

- [1] Cordero P and Salamó S 1993 *Found. Phys.* **23** 675 ; 1994 *J. Math. Phys.* **35** 3301
- [2] Natanzon G A 1979 *Teor. Mat. Fiz.* **38** 146
- [3] del Sol Mesa A, Quesne C and Smirnov Y F 1998 *J. Phys. A: Math. Gen.* **31** 321
- [4] For a complete review of SUSYQM see Cooper F, Khare A and Sukhatme U 1995 *Phys. Rep.* **251** 2671
- [5] Gendenshtein L 1983 *JETP Lett.* **38** 356
- [6] Codriansky S, Cordero P and Salamó 1999 S *J. Phys. A: Math. Gen.* **32** 6287
- [7] Gradshteyn I S and Ryzhik I M 1965 *Table of Integrals, Series and Products* (New York: Academic Press)
- [8] Cooper F, Ginocchio J N and Khare A 1987 *Phys. Rev. D* **36** 2458

- [9] Dabrowska J, Khare A and Sukhatme A P 1988 *J. Phys. A: Math. Gen.* **21** L195
- [10] Alhassid Y, Gürsey F and Iachello F 1986 *Ann Phys* **167** 181